

ONE-SIDED IDEALS AND APPROXIMATE IDENTITIES IN OPERATOR ALGEBRAS

DAVID P. BLECHER*

ABSTRACT. A left ideal of any C^* -algebra is an example of an operator algebra with a right contractive approximate identity (r.c.a.i.). Indeed left ideals in C^* -algebras may be characterized as the class of such operator algebras, which happen also to be triple systems. Conversely, we show here and in a sequel to this paper [9], that operator algebras with r.c.a.i. should be studied in terms of a certain left ideal of a C^* -algebra. We study left ideals from the perspective of ‘Hamana theory’ and using the multiplier algebras introduced by the author. More generally, we develop some general theory for operator algebras which have a 1-sided identity or approximate identity, including a Banach-Stone theorem for these algebras, and an analysis of the ‘multiplier operator algebra’.

*This research was supported by a grant from the National Science Foundation.

*Many of these results were presented at the November '01 Meeting of the A.M.S. in Irvine, California. Sept 14, 2001. Revision of November 12, '01.

1. INTRODUCTION AND NOTATION

A left ideal of any C^* -algebra is an example of an operator algebra with a right contractive approximate identity. Conversely, we shall see here, and in a sequel to this paper [9], that operator algebras with a right contractive approximate identity should be studied in terms of a certain left ideal of a C^* -algebra.

A (concrete) operator algebra is a subalgebra of $B(H)$, for some Hilbert space H . More abstractly, an operator algebra will be an algebra A with a norm defined on the space $M_n(A)$ of $n \times n$ matrices with entries in A , for each $n \in \mathbb{N}$, such that there exists a *completely isometric*¹ homomorphism $A \rightarrow B(H)$ for some Hilbert space H . In this paper all our operator algebras and spaces will be taken to be complete. We shall say that an operator algebra is *unital* if it has a two-sided contractive identity. Unital operator algebras were characterized abstractly in [12]. However the class of nonselfadjoint operator algebras which is of most interest to C^* -algebraists or those interested in noncommutative geometry is the class of left or right ideals in a C^* -algebra, which is a nonselfadjoint operator algebra with a right (respectively left) contractive approximate identity. Thus in this paper we shall study operator algebras with a one-sided (usually right) contractive approximate identity. We shall abbreviate ‘right (resp. left) contractive approximate identity’ to ‘r.c.a.i.’ (resp. ‘l.c.a.i.’). It is well known even for Banach algebras that an algebra with both a left and a right c.a.i. has a two-sided c.a.i..

As we point out in §2, left ideals in C^* -algebras may be characterized as the class of nonselfadjoint operator algebras with r.c.a.i., which happen also to be ‘triple systems’. This therefore suggests that left ideals in C^* -algebras may profitably be studied using machinery that exploits both the ‘operator algebra’ and the ‘triple’ structure, and indeed we do take this approach here. For example ‘morphisms’ of left ideals will be what we call ‘ideal homomorphisms’ below, namely homomorphisms which are also ‘triple morphisms’.

Much of this paper is concerned with left ideals in unital operator algebras which have a r.c.a.i.. In a formal sense this coincides with the class of all operator algebras with r.c.a.i., since every operator algebra is easily seen to be a left ideal in some operator algebra. By Proposition 6.4 in [8] (although we shall not use this fact except for motivational purposes), this also coincides with the class of left M-ideals in unital operator algebras. We shall not however use this last fact, except as evidence that these objects are more important than they seem at first. We will make the blanket convention in this paper that all ideals, left or otherwise, are closed, i.e. complete.

We are not aware of any general results in the literature on operator algebras with r.c.a.i., and one of the main goals of this paper and its sequel [9] is to take on such a study. Although operator algebras with r.c.a.i. have not hitherto come up in the literature on general operator algebras as much as their two-sided relative, they do arise very naturally (e.g. left ideals in a C^* -algebra, Ae for a projection e in a unital operator algebra A ; or Example 2.4, or Theorem 4.11 in [10]). We also felt that it was worthwhile to dispel the common feeling that only operator algebras with 2-sided approximate identities have any satisfactory theory. Indeed we show amongst other things that they have an abstract characterization, Banach-Stone type theorems, reasonable multiplier algebras (which are operator algebras with two sided

¹A map $T : X \rightarrow Y$ is completely isometric if $[x_{ij}] \mapsto [T(x_{ij})]$ is isometric on $M_n(X)$ for all $n \in \mathbb{N}$.

identity of norm 1), and they have an operator space predual if and only if they are dual operator algebras' in the usual strong sense of that term. Also, this subject becomes a little more interesting with a certain 'transference principle' in mind. This principle (which is proved fully in the sequel [9]), allows one to deduce many general results about operator algebras with r.c.a.i., from results about left ideals in a C^* -algebra. Namely there is an important left ideal $\mathfrak{J}_e(A)$ of a C^* -algebra $\mathcal{E}(A)$, which is associated to any such operator algebra. We call $\mathfrak{J}_e(A)$ the 'left ideal envelope' of A . This is analogous to what happens in the case of operator algebras with 2-sided c.a.i., which are largely studied these days in terms of a certain C^* -algebra, namely the C^* -envelope.

We now describe the layout of the paper. In §2 we give many preliminary lemmas which will be used later and in [9], as well as many new results about left ideals and operator algebras with r.c.a.i..

In §3 we look at Banach-Stone type theorems. The classical Banach-Stone theorem (see e.g. [14] IV.2) may be stated in the following form: if $C(K_1) \cong C(K_2)$ linearly isometrically, then they are $*$ -isomorphic (from which it is clear that the compact spaces K_1 and K_2 are homeomorphic). Indeed the usual proofs show that the linear isometry equals a $*$ -isomorphism $C(K_1) \rightarrow C(K_2)$ multiplied by a fixed unitary in $C(K_2)$. Note that the linear isometry is unital (i.e. takes the $1 \mapsto 1$) if and only if $u = 1$, and one often proves this special case first. There are numerous noncommutative versions of this, the most well known due to Kadison [22], where the $C(K)$ spaces are replaced by C^* -algebras. For unital maps between nonselfadjoint operator algebras with two-sided identities of norm 1, a similar theorem is true, that is, any unital linear complete isometry between such operator algebras, is also a homomorphism. This is an immediate consequence of the existence of the Arveson-Hamana C^* -envelope of [1, 2, 20] (indeed this generalization of the Banach-Stone theorem was a major consideration in [1, 2], see also [3, 16]). In [6] Appendix B.1 it is shown that a linear surjective complete isometry $\varphi : A \rightarrow B$ between operator algebras with contractive approximate identities may be written as $\varphi = \pi(\cdot)u$ for a surjective completely isometric homomorphism $\pi : A \rightarrow B$, and a unitary u with $u, u^* \in M(B)$. Here $M(B)$ is the multiplier algebra (see [31] for example). In §3 below we examine such theorems in the case of operator algebras with one-sided identities or contractive approximate identities, for example in the case of left ideals in a C^* -algebra. We see that the Murray-von Neumann equivalence of projections plays a role here.

In §4 we study the 'left multiplier operator algebra' $LM(A)$ of an operator algebra A with l.c.a.i. (which will be a symmetrical theory to that of $RM(A)$ for an operator algebra A with r.c.a.i.). In the next paper we develop a satisfactory candidate for $LM(A)$ if A has a r.c.a.i. - this does not work out quite as nicely as the case considered in §4, unless A is a left ideal of a C^* -algebra.

We remark that we are not aware even of much general theory of one-sided contractive approximate identities in general Banach algebras. The only references we can give for this subject are the works of P. G. Dixon (see [26] for references), and the general texts [13, 26].

Often it is convenient to state only the 'right-handed' version of a result, say. For example Lemma 2.8 is a result about operator algebras with r.c.a.i.. Of course by symmetry there will be a matching 'left-handed' version, in our example it will be about operator algebras

with l.c.a.i.. If we want to invoke this ‘left-handed’ version, we will refer to the ‘other-handed version of Lemma 2.8’, for example. Sometimes we will be sloppy and simply refer to Lemma 2.8, and leave it to the reader to check that the result referred to does indeed have a symmetric or ‘other-handed version’.

We end the introduction with some more notation, and some background results which will be useful in various places. We reserve the letters H, K for Hilbert spaces, and J for a left ideal of a C^* -algebra.

If A is an algebra then we write $\lambda : A \rightarrow \text{Lin}(A)$ for the canonical ‘left regular representation’ of A on itself. If S is a subalgebra of A then the left idealizer of S is the subalgebra $\{x \in A : xS \subset S\}$ of A . Note S is a left ideal in this subalgebra, whence the name. Similarly for the right idealizer; the (2-sided) idealizer is the intersection of the left and right idealizer.

By a ‘representation’ $\pi : A \rightarrow B(H)$ of an operator algebra A we shall mean a completely contractive homomorphism. If A has r.c.a.i. and if we say that π is *nondegenerate*, then at the very least we mean that $\pi(A)H$ is dense in H . Note that this last condition does not imply in general that $\pi(e_\alpha)\zeta \rightarrow \zeta$ for $\zeta \in H$, where $\{e_\alpha\}$ is the r.c.a.i., as one is used to in the two-sided case. One also cannot appeal to Cohen’s factorization theorem in its usual form (see however [26] §5.2).

We will use without comment several very basic facts from C^* -algebra theory (see e.g. [29]), such as the basic definitions of the left multiplier algebra $LM(A)$ of a C^* -algebra, and the multiplier algebra $M(A)$.

As a general reference for operator spaces the reader might consult [17, 30] or [27]. We write $\hat{} : X \rightarrow X^{**}$ for the canonical map, this is a complete isometry if X is an operator space, and is a homomorphism if X is an operator algebra (giving the second dual the Arens product [13]).

We will often consider the basic examples C_n (resp. R_n) of operator algebras with right (resp. left) identity of norm 1; namely the $n \times n$ matrices ‘supported on’ the first column (resp. row). This is a left (resp. right) ideal of M_n , and has the projection E_{11} as the 1-sided identity. We write $C_n(X)$ for the first column on $M_n(X)$, that is $M_{n,1}(X)$. If X is an operator space so is $C_n(X)$.

If X and Y are subsets of an operator algebra we usually write XY for the *norm closure* of the set of finite sums of products xy of a term in X and a term in Y . For example, if J is a left ideal of a C^* -algebra A , then with this convention J^*J will be a norm closed C^* -algebra. This convention extends to three sets, thus $JJ^*J = J$ for a left ideal of a C^* -algebra as is well known (or use the proof of Lemma 2.1 below to see this).

We recall more generally that a TRO (*ternary ring of operators*) is a (norm closed for this paper) subspace X of $B(K, H)$ such that $XX^*X \subset X$. It is well known (copy the proof of Lemma 2.1 below) that in this case $XX^*X = X$. Then XX^* and X^*X are C^* -algebras, which we will call the left and right C^* -algebras of X respectively, and X is a $(XX^*) - (X^*X)$ -bimodule. It is also well known that TRO’s are the same thing as Hilbert C^* -modules. A linear map $T : X \rightarrow Y$ between TRO’s is a *triple morphism* if $T(xy^*z) = T(x)T(y)^*T(z)$ for all $x, y, z \in X$. TRO’s are operator spaces, and triple morphisms are completely contractive, and indeed are completely isometric if they are 1-1 (see e.g. [21], this is related to results of Harris and Kaup). A completely isometric surjection between

TRO's is a triple morphism. This last result might date back to around 1986, to Hamana and Ruan's PhD thesis independently. See [21] or [6] A.5 for a proof.

If J is a left ideal in a C^* -algebra, and if B is a C^* -algebra, then a certain class of maps $J \rightarrow B$ are of particular importance in this paper. Namely, those which are restrictions of $*$ -representations $\theta : JJ^* \rightarrow B(H)$. We will call these *ideal homomorphisms* of J . They are characterized in 2.10 below as the homomorphisms which are also triple morphisms.

Next we recall the left multiplier algebra $\mathcal{M}_\ell(X)$ of an operator space X . This is a unital operator algebra, which may be viewed as a subalgebra of $CB(X)$ containing Id_X , but with a different (bigger in general) norm. Here $CB(X)$ are the 'completely bounded' linear maps on X . One may take the definition of $\mathcal{M}_\ell(X)$ from the following result from [8]:

Theorem 1.1. *A linear $T : X \rightarrow X$ on an operator space is in $Ball(\mathcal{M}_\ell(X))$ if and only if $T \oplus Id$ is a complete contraction $C_2(X) \rightarrow C_2(X)$.*

As we said above, $\mathcal{M}_\ell(X)$ is an operator algebra. The matrix norms on $\mathcal{M}_\ell(X)$ may be described via the natural isomorphism $M_n(\mathcal{M}_\ell(X)) \cong \mathcal{M}_\ell(M_n(X))$. That is the norm of a matrix $[T_{ij}]$ of multipliers may be taken to be the norm in $\mathcal{M}_\ell(M_n(X))$ of the map

$$[x_{ij}] \mapsto \left[\sum_k T_{ik}(x_{kj}) \right] .$$

There are several other equivalent definitions of $\mathcal{M}_\ell(X)$ given in [6, 8, 11]. One of our main motivations for the introduction of multipliers of operator spaces in [6] was in order to give a more unified and 'extremal' approach to the main theorems characterizing operator algebras and modules. We pointed out in §5 of [6] that in order to prove the characterization of operator algebras of [8] say, it is clearly only necessary to check that the 'left regular representation' $\lambda : A \rightarrow CB(A)$, is a complete isometry into the operator algebra $\mathcal{M}_\ell(A)$. But this is immediate from a theorem such as 1.1 above - see the simple proof of Lemma 2.13 for more details if required. (This deduction of [8] from 1.1 was noticed independently by Paulsen, who also gave a simplified proof of 1.1). The same proof works to give immediately the following characterization of operator algebras with one-sided c.a.i., which is a slight variation on [7] 1.11 to which the reader is referred if more details of proof are needed (we warn the reader that the statement of that result is stated slightly incorrectly, it neglected to mention that the matrices in A there should have norm ≤ 1).

Theorem 1.2. *Let A be an operator space which is an algebra with a right identity of norm 1 or r.c.a.i.. Then A is completely isometrically isomorphic to a concrete operator algebra (via a homomorphism of course), if and only if we have*

$$\|(x \oplus Id_n)y\| \leq 1$$

for all $n \in \mathbb{N}$ and $x \in Ball(M_n(A)), y \in Ball(M_{2n,n}(A))$.

To explain the notation of the theorem, we have written Id for a formal identity, thus the expression $(x \oplus Id_n)y$ above means that the upper $n \times n$ -submatrix of y is left multiplied by x , and the lower submatrix is left alone.

Acknowledgements: We thank M. Kaneda for catching several misprints, and for several conversations related to the present paper and [9].

2. GENERAL RESULTS AND LEMMATA

In this section we collect many simple lemmas and other background facts which will be used later and in [9]. Some of the results here for left ideals in C^* -algebras may be known, but in any case our proofs are short. Certainly the first lemma is classical and extremely well known; we have appended a (well known) proof which might be convenient for the reader who accepts the existence of c.a.i.'s in C^* -algebra (modern texts in this area contain short proofs of this latter fact (see e.g. [19])).

Lemma 2.1. *(Classical) A (norm closed) left ideal J in a C^* -algebra is an operator algebra with a positive right contractive approximate identity. Also $J \cap J^* = J^*J \subset J \subset JJ^*$, so that J is a left ideal of JJ^* .*

Proof. A left ideal J in a C^* -algebra A is clearly a subalgebra of A . Also JJ^* and J^*J are C^* -subalgebras of A . So J^*J has a positive c.a.i. $\{e_\alpha\}$. Then for $x \in J$,

$$\|xe_\alpha - x\|^2 = \|e_\alpha x^* x e_\alpha - x^* x e_\alpha - e_\alpha x^* x + x^* x\| \rightarrow 0.$$

The remaining assertions follow immediately from this; for example if $x \in J \cap J^*$ then $x^* = \lim x^* e_\alpha$, so that $x \in J^*J$. $\square$

Note that J^*J also equals $\{x \in J : e_\alpha x \rightarrow x\}$, where $\{e_\alpha\}$ is the c.a.i. for J mentioned above. This is part of our motivation for the next definition and result (which will only be used in part of §4)².

Definition 2.2. *We say that an operator algebra A with r.c.a.i. (resp. l.c.a.i.) has property $(\mathcal{R})$ (resp. $(\mathcal{L})$) if a r.c.a.i. (resp. l.c.a.i.) $\{e_\alpha\}$ exists for A such that $e_\alpha e_{\alpha'} \rightarrow e_{\alpha'}$ (resp. $e_{\alpha'} e_\alpha \rightarrow e_{\alpha'}$) for each fixed $e_{\alpha'}$ in the net. In this case we define $\mathcal{R}(A) = \{x \in A : e_\alpha x \rightarrow x\}$ (resp. $\mathcal{L}(A) = \{x \in A : x e_\alpha \rightarrow x\}$).*

Remarks. We note that a left ideal of a C^* -algebra has property $(\mathcal{R})$, and in this case $\mathcal{R}(A) = J^*J$. More generally a subalgebra of a C^* -algebra with a self-adjoint right c.a.i. has property $(\mathcal{R})$, since in this case $(e_\alpha e_{\alpha'})^* = e_{\alpha'} e_\alpha \rightarrow e_{\alpha'} = e_{\alpha'}^*$. An operator algebra with two-sided c.a.i. obviously has property $(\mathcal{R})$, and in this case $\mathcal{R}(A) = A$. Certainly every operator algebra with a right identity of norm 1 has property $(\mathcal{R})$. We are not aware of any operator algebras with r.c.a.i. which do not have property $(\mathcal{R})$ (if all do this would certainly solve a problem encountered in [10] §4 and elsewhere in some of the authors work: whether every ‘Hilbertian module’ is a rigged module). Thus it seems a reasonable and general property.

Proposition 2.3. *If an operator algebra A with r.c.a.i. has property $(\mathcal{R})$, then $\mathcal{R}(A)$ is a norm closed right ideal of A (and hence is an operator algebra) with two sided c.a.i.. Moreover, $\mathcal{R}(A)$ does not depend on the particular c.a.i. $\{e_\alpha\}$ considered. Also, $A \mathcal{R}(A) = A$ and $\mathcal{R}(A) A = \mathcal{R}(A)$.*

Similar results hold for property $(\mathcal{L})$.

Proof. The first assertion we leave as a simple exercise. Suppose that A has property $(\mathcal{R})$ with respect to one r.c.a.i. $\{e_\alpha\}$, and let $\{f_\beta\}$ be another r.c.a.i. such that $f_\beta f_{\beta'} \rightarrow f_{\beta'}$ for every fixed β' . Let $B = \{a \in A : f_\beta a \rightarrow a\}$, another right ideal of A with two sided

²Another part of our motivation and inspiration is the theory in [4] and §4 of [10].

c.a.i.. Note that $\mathcal{R}(A)B = \mathcal{R}(A)$ and $B\mathcal{R}(A) = B$. Thus by (the other-handed version of) [10] Theorem 4.15, $B = \mathcal{R}(A)$. The remaining assertions are left to the reader. $\square$

Example 2.4.

Let B be a unital operator algebra, a unital subalgebra of a W^* -algebra N , and define $M_\infty(N)$ to be the von Neumann algebra $B(\ell^2) \bar{\otimes} N$, thought of as infinite matrices $[b_{ij}]$ with entries b_{ij} indexed over $i, j \in \mathbb{N}$. We let $M_\infty(B)$ be the subset of $M_\infty(N)$ consisting of those matrices with entries b_{ij} in B . Let $C_\infty^w(B)$ be the ‘first column’ of $M_\infty(B)$, and let $R_\infty(B)$ be the space of row vectors $[b_1 b_2 \cdots]$ with entries $b_i \in B$, such that $\sum_k b_k b_k^*$ converges in norm. These spaces are familiar objects in operator space theory (see [17]), and also in C^* -module theory. We may then consider the closed subspace $A = C_\infty^w(B)R_\infty(B)$ of $M_\infty(B)$; those familiar with operator space theory will have no trouble verifying that A is a subalgebra of $M_\infty(N)$, that A has a nonnegative r.c.a.i., and indeed if $B = N$ then A is a left ideal of $M_\infty(B)$. In fact A contains the C^* -algebra $\mathbb{K}_\infty(B)$, namely the spatial tensor product $\mathbb{K}(\ell_2) \otimes B$ (which in the language of C^* -modules equals $\mathbb{K}(C_\infty(A))$), and the usual c.a.i. for this C^* -algebra, namely $I_n \otimes 1_B$, is a r.c.a.i. for A . Thus A has property $(\mathcal{R})$. It is easily verified that $\mathbb{K}_\infty(B)$ is a right ideal in A , and in fact $\mathcal{R}(A) = \mathbb{K}_\infty(B)$. This algebra A is important to those working on the borders of operator spaces and operator algebras, although it has rarely appeared in the literature as far as we are aware, and that fleetingly.

The next lemma concerns ‘principal ideals’. By a ‘principal ideal’ in a C^* -algebra A , we will mean by analogy with pure algebra, an ideal of the form Ax for some $x \in A$. We are not taking the norm closure here, $Ax = \{ax : a \in A\}$ for some $x \in A$; however in view of the importance of closed ideals in C^* -algebra theory, below we will only consider principal ideals which are already norm closed. We remark however that if A is nonunital, then when transferring results of the kind found in pure ring theory to the C^* -algebra case, one finds that the multiplier algebra of A plays an important role. Hence the reader might argue that ideals of the form Ax with $x \in M(A)$ (or $RM(A)$), should be thought of as ‘principal ideals’ of A too. However in our paper we shall not need these more general ideals, and so stick with the earlier definition for convenience. Interestingly though, all these kind of ideals have a simplified form:

Proposition 2.5. *Let A be a C^* -algebra, and suppose that $J = Ax$ is a closed left ideal, with $x \in A$ (resp. $x \in M(A)$). Then $J = Ae$ where e is an orthogonal projection in J (resp. in $M(A)$).*

Proof. Since J is the range of an adjointable map on A , J is orthogonally complemented in the sense of C^* -module theory, by [32] 15.3.9. This implies that $J = Ae$ where e is an orthogonal projection in $M(A)$. So that if A is unital we are done, and note that in this case Ae has a right identity of norm 1. However in any case, if $x \in A$, then $Ax = M(A)x$ (clearly $Ax \subset M(A)x$, but if $T \in M(A)$ then $Tx = \lim T e_\alpha x \in Ax$). Thus applying the above we see that J has a right identity f of norm 1, and $f \in J \subset A$. Hence $J = Af$. $\square$

Corollary 2.6. *Suppose that J is a left ideal of a C^* -algebra, and suppose that J has a right identity. Then J also has a right identity of norm 1. Moreover the latter is the norm limit of any r.c.a.i. for the left ideal.*

Proof. Let J be the left ideal, which by the previous result, has a right identity e of norm 1. So $e = e^* \in J \cap J^* = J^*J$. If $\{e_\alpha\}$ is a r.c.a.i. for J then $\{e_\alpha^*e_\alpha\}$ is a 2-sided c.a.i. for J^*J (see e.g. 2.9), thus $e_\alpha^*e_\alpha = e_\alpha^*e_\alpha e \rightarrow e$. Finally,

$$\|e_\alpha - e\|^2 = \|e_\alpha^*e_\alpha - e_\alpha^*e - ee_\alpha + e\| \rightarrow 0.$$

□

Later we will prove the analogous result to the last corollary valid for operator algebras.

Lemma 2.7. *Suppose that A is an operator algebra with two right identities e and f of norm 1. Then $e = f$.*

Proof. Since e and f are orthogonal projections we have $e = ef = e^* = fe = f$. □

Lemma 2.8. ([13] 28.7). *If A is an operator algebra with r.c.a.i. then A^{**} is an operator algebra with right identity of norm 1. If A has a right identity e , then $\hat{e}$ is the right identity of A^{**} .*

The last assertion of the following lemma is well known; however we state it here because it is a trivial corollary of the more important (for us) first part:

Lemma 2.9. *Suppose that $a \in B(H)$, and $\{e_\alpha\}$ is a net of contractions in $B(H)$ such that $ae_\alpha \rightarrow a$. Then $ae_\alpha e_\alpha^* \rightarrow a$, $ae_\alpha^* e_\alpha \rightarrow a$, and $ae_\alpha^* \rightarrow a$.*

Thus if J is a left ideal of a C^ -algebra, and if $\{e_\alpha\}$ is a r.c.a.i. for J , then $\{e_\alpha^*e_\alpha\}$ is a nonnegative right contractive approximate identity for J (and indeed also is a 2-sided c.a.i. for the C^* -subalgebra $J \cap J^* = J^*J$).*

Proof. We use a technique from [10]. If $ae_\alpha \rightarrow a$ then $ae_\alpha e_\alpha^* a^* \rightarrow aa^*$, so that $0 \leq a(I - e_\alpha e_\alpha^*)a^* \rightarrow 0$. Thus by the C^* -identity, $a\sqrt{I - e_\alpha e_\alpha^*} \rightarrow 0$. Multiplying by $\sqrt{I - e_\alpha e_\alpha^*}$ we see that $a(I - e_\alpha e_\alpha^*) \rightarrow 0$ as required for the first assertion. Also,

$$\|ae_\alpha^* - a\| \leq \|ae_\alpha^* - ae_\alpha e_\alpha^*\| + \|ae_\alpha e_\alpha^* - a\| \rightarrow 0$$

since $\|ae_\alpha^* - ae_\alpha e_\alpha^*\| \leq \|a - ae_\alpha\| \rightarrow 0$. Finally,

$$\|ae_\alpha^* e_\alpha - a\| \leq \|ae_\alpha^* e_\alpha - ae_\alpha\| + \|ae_\alpha - a\| \leq \|ae_\alpha^* - a\| + \|ae_\alpha - a\| \rightarrow 0$$

by what we just proved. □

Note that the last result shows that a r.c.a.i. for a C^* -algebra is a l.c.a.i. too.

If J is a left ideal in a C^* -algebra, then we recall from §1 that an *ideal representation* or *ideal homomorphism* of J is a restriction of a $*$ -representation $\theta : JJ^* \rightarrow B(H)$ to J . Clearly such a map is completely contractive.

Proposition 2.10. *Let J be a left ideal of a C^* -algebra, and let $\pi : J \rightarrow B(H)$ be a function. Then π is the restriction of a $*$ -representation $\theta : JJ^* \rightarrow B(H)$ if and only if π is a homomorphism and a triple morphism. Moreover such π is completely isometric if and only if π is 1-1, and if and only if θ is 1-1.*

Proof. If π is the restriction of a $*$ -representation then it is evident that π is a homomorphism and a triple morphism. Conversely, it is well known (see [21]), that if π is a triple morphism, then there is an associated $*$ -homomorphism $\theta : JJ^* \rightarrow B(H)$ with the property that $\theta(xy^*) = \pi(x)\pi(y)^*$ for all $x, y \in J$. If in addition π is a homomorphism, and $\{e_\alpha\}$ is a positive r.c.a.i. for J , then $\{\pi(e_\alpha)\}$ is a positive r.c.a.i. for $\pi(J)$, and so for $x \in J$,

$$\theta(x) = \lim \theta(xe_\alpha) = \lim \pi(x)\pi(e_\alpha)^* = \pi(x)$$

by Lemma 2.9.

If further π is 1-1, then it is shown in [21] that θ is 1-1. $\square$

The following result is a simple consequence of the fact that $JJ^*J = J$:

Lemma 2.11. *Let J be a left ideal of a C^* -algebra, and let $\theta : JJ^* \rightarrow B(H)$ be a $*$ -homomorphism. If π is the restriction to J then θ is nondegenerate if and only if $\pi(J)H$ is dense in H .*

If A has a left identity of norm 1 but no right identity, and if $\pi : A \rightarrow B(H)$ is a nondegenerate isometric representation, then $\pi(e) = Id$, so that $\pi(ae) = \pi(a)$, so that $ae = a$ for all $a \in A$. This is a contradiction. Thus there is in general little point in seeking nondegenerate isometric representations of algebras with l.c.a.i.. (The reader may think at this juncture of ‘adjoint nondegeneracy’ but this really is a different issue to the point we are now making). The following discussion, proposition, and subsequent definition of $\mathcal{L}$ -isometric representation, gives one way to fix the above problem.

If A has left identity e of norm 1, then A clearly has property $(\mathcal{L})$ of 2.2, and this identity is the 2-sided identity of $\mathcal{L}(A) = Ae$. Moreover, the map $A \rightarrow \mathcal{L}(A)$ taking $a \mapsto ae$, is a completely contractive homomorphism, and also is a complete quotient map and indeed is a projection onto $\mathcal{L}(A)$. On the other hand, if A has a l.c.a.i. and property $(\mathcal{L})$, then we can make similar assertions for the second dual using 2.8: there is a completely contractive homomorphism, which is a complete quotient map and indeed a projection $A^{**} \rightarrow \mathcal{L}(A)^{**}$. This is the map $F \mapsto FE$, where E is a weak* limit point of the c.a.i. of $\mathcal{L}(A)$.

Proposition 2.12. *Suppose that A is an operator algebra with l.c.a.i. and property $(\mathcal{L})$ of 2.2. Let $\pi : A \rightarrow B(H)$ be a completely contractive representation (resp. and also $\pi(A)H$ is dense in H). Then $\pi|_{\mathcal{L}(A)} : \mathcal{L}(A) \rightarrow B(H)$ is a completely contractive homomorphism (resp. and also such that $\pi(\mathcal{L}(A))H$ is dense in H). Conversely, if $\theta : \mathcal{L}(A) \rightarrow B(H)$ is a completely contractive homomorphism, then there exists a completely contractive homomorphism $\pi : A \rightarrow B(H)$ extending θ . If further $\theta(A)H$ is dense in H then π is unique, and $\pi(A)H$ is dense in H . Finally,*

$$\{T \in B(H) : T\pi(A) \subset \pi(A)\} = \{T \in B(H) : T\pi(\mathcal{L}(A)) \subset \pi(\mathcal{L}(A))\}.$$

Proof. The first statements are simple exercises. For the converse, given such $\theta : \mathcal{L}(A) \rightarrow B(H)$, consider the series of completely contractive homomorphisms

$$A \hookrightarrow A^{**} \rightarrow \mathcal{L}(A)^{**} \xrightarrow{\theta^{**}} B(H)^{**} \rightarrow B(H).$$

The homomorphism $A^{**} \rightarrow \mathcal{L}(A)^{**}$ is the map described above the Proposition, and the other maps are the canonical ones. The composition of these homomorphisms is the desired

π . We leave it to the reader to check the details. Since $\pi(a)\theta(b)\zeta = \pi(ab)\zeta = \theta(ab)\zeta$ for $a \in A, b \in \mathcal{L}(A), \zeta \in H$ we see that π is unique if $\pi(A)H$ is dense.

Finally, using the ‘other-handed version’ of the last assertion of 2.3, we see for example that if $T\pi(A) \subset \pi(A)$ then $T\pi(\mathcal{L}(A)) = T\pi(A)\pi(\mathcal{L}(A)) \subset \pi(A)\pi(\mathcal{L}(A)) = \pi(\mathcal{L}(A))$. The other direction is similar. $\square$

The previous result shows that A and $\mathcal{L}(A)$ have the same representation theory. Thus the following definition which plays a role in §4 is somewhat natural: we say that a nondegenerate representation $\pi : A \rightarrow B(H)$ is completely ‘ $\mathcal{L}$ -isometric’, if $\pi|_{\mathcal{L}(A)}$ is completely isometric on $\mathcal{L}(A)$.

Lemma 2.13. *Let A be an operator algebra with a r.c.a.i.. Then the canonical ‘left regular representation’ of A on itself yields completely contractive embeddings (i.e. 1-1 homomorphisms)*

$$A \hookrightarrow \mathcal{M}_\ell(A) \hookrightarrow CB(A) ,$$

and the first of these embeddings, and their composition, are completely isometric.

Proof. Let $\lambda : A \rightarrow CB(A)$ be the left regular representation. This map is certainly completely contractive, however since $\lambda(a)(e_\alpha) = ae_\alpha \rightarrow a$ it is clear that it is a complete isometry. It is also clear that if $a \in \text{Ball}(A)$ then $\lambda(a)$ satisfies the criterion of 1.1, so that $\lambda(a) \in \text{Ball}(\mathcal{M}_\ell(A))$. A similar argument works at the matrix level. Thus λ factors through $\mathcal{M}_\ell(A)$ via the two completely contractive homomorphisms above. Since λ is completely isometric, so is the first embedding. $\square$

We remark that the canonical embedding $\mathcal{M}_\ell(A) \hookrightarrow CB(A)$, where A is an operator algebra with r.c.a.i., is not in general completely isometric, or even isometric (an example of this is given in [9]). This has implications for our theory of multipliers in [9].

The following ‘Hamana-theory’ type results, which are very interesting in their own right, will be one of our main tools to deduce results about operator algebras with r.c.a.i., from results about left ideals in a C^* -algebra. We give condensed versions of two of these results from [9].

We say that a pair (J, i) consisting of a left ideal J in a C^* -algebra, and a completely isometric homomorphism $i : A \rightarrow J$, is a *left ideal extension* of A if $i(A)$ ‘generates J as a TRO’. That is, the span of expressions of the form $i(a_1)i(a_2)^*i(a_3)i(a_4)^*\cdots i(a_{2n+1})$, for $a_i \in A$, are dense in J . It follows from this that $\{i(e_\alpha)\}$ is a r.c.a.i. for J if $\{e_\alpha\}$ is a r.c.a.i. for A .

Theorem 2.14. *Let A be an operator algebra with r.c.a.i.. Then there exists a left ideal extension $(\mathfrak{J}_e(A), j)$ of A , with $\mathfrak{J}_e(A)$ a left ideal in a C^* -algebra $\mathcal{E}(A)$, such that for any other left ideal extension (J, i) of A , there exists a (necessarily unique and surjective) ideal homomorphism $\tau : J \rightarrow \mathfrak{J}_e(A)$ such that $\tau \circ i = j$. Thus $\mathfrak{J}_e(A)/(\ker \tau) \cong J$ completely isometrically homomorphically (i.e as operator algebras) too. Moreover $(\mathfrak{J}_e(A), j)$ is unique in the following sense: given any other (J', j') with this universal property, then there exists a surjective completely isometric homomorphism $\theta : \mathfrak{J}_e(A) \rightarrow J'$ such that $\theta \circ j = j'$.*

Finally, $(\mathfrak{J}_e(A), j)$ is a triple envelope for A in the sense of [21].

We call $(\mathfrak{J}_e(A), j)$ the *left ideal envelope* of A , and often write $\mathcal{E}(A) = \mathfrak{J}_e(A)\mathfrak{J}_e(A)^*$. The map j will be called the *Shilov embedding homomorphism*. From the last assertion of the theorem, and the first definition of $\mathcal{M}_\ell(A)$ given in §4 of [6], we may identify $\mathcal{M}_\ell(A)$ with $\{R \in LM(\mathcal{E}(A)) : Rj(A) \subset j(A)\}$. If $\mathcal{E}(A)$ is represented nondegenerately as a C^* -subalgebra of $B(H)$, then we can also identify $\mathcal{M}_\ell(A)$ with $\{R \in B(H) : Rj(A) \subset j(A)\}$, completely isometrically isomorphically. We shall not however use this last remark.

Corollary 2.15. *Let A be an operator algebra with r.c.a.i., and λ the usual left regular representation of A . Then for any $T \in \mathcal{M}_\ell(A)$, regarded as a map on A , we have $T\lambda(A) \subset \lambda(A)$. Thus elements of $\mathcal{M}_\ell(A)$, considered as maps on A , are right A -module maps. That is, $\mathcal{M}_\ell(A) \subset CB_A(A)$ as sets. Also, $\mathcal{M}_r(A) \subset {}_A CB(A)$ as sets.*

Proof. The first assertion follows from the penultimate line before the statement of the Corollary, together with the fact that j is a homomorphism. For if $a \in A$, then the map $b \mapsto ab$ on A , corresponds to the map $j(b) \mapsto j(a)j(b)$ on $j(A)$. Thus if the left multiplier T corresponds to an $R \in LM(\mathcal{E}(A))$ with $Rj(a) = j(T(a))$ then $Rj(ab) = j(T(a)b)$ for any $b \in A$, which amounts to the first assertion, and also yields the second assertion immediately. The third is similar. $\square$

Theorem 2.16. ([9]) *Let A be an operator algebra with a r.c.a.i.. Then there exists an injective unital C^* -algebra B containing $\mathcal{E}(A)$ as a C^* -subalgebra, and an orthogonal projection $e \in B$, such that (Be, j) is an injective envelope for A , where j is the Shilov embedding homomorphism. Moreover if A has a right identity f then $j(f) = e$.*

We define the *ideal injective envelope* of A to be the ideal Be in the last theorem.

Corollary 2.17. *Let J be a left ideal in a C^* -algebra. Then the canonical embedding of a left ideal in a C^* -algebra, into its ideal injective envelope Be is an ideal homomorphism.*

Proof. Using 2.14, there exists a surjective ideal homomorphism $\tau : J \rightarrow \mathfrak{J}_e(J)$, such that $\tau = j$. Clearly $\mathfrak{J}_e(J)$ is a subalgebra and sub-TRO of $\mathcal{E}(J)$, and consequently also of B and Be . $\square$

As another corollary of [9], one may give a characterization of left ideals in C^* -algebras. We will say that an operator space X is an abstract triple system if it is linearly completely isometrically isomorphic to a TRO Z . Note that then one may pull back the triple product on Z to a triple product $\{\cdot, \cdot, \cdot\}$ on X , and by the TRO result of Harris and Ruan mentioned in the introduction, this triple product on X is unique, i.e. independent of Z . That is, this triple product is completely determined by the ‘operator space structure’ or matrix norms on X . Indeed Neal and Russo have a striking recent characterization of abstract triple systems in terms of these matrix norms [25]. Putting this together with the theorem below, and a characterization of operator algebras with right contractive approximate identity (r.c.a.i.) which we gave in Theorem 1.2 for example, gives a ‘completely abstract’ characterization of left ideals in C^* -algebras.

Theorem 2.18. *Let A be an abstract operator algebra which is also an abstract triple system (we are assuming the underlying matrix norms for both structures coincide). The following are equivalent:*

- (i) *A has a r.c.a.i. for the algebra product.*
- (ii) *A has a r.c.a.i. $\{e_\alpha\}$ for the algebra product such that $\{a, e_\alpha, b\} \rightarrow ab$ for all $a, b \in A$;*
- (iii) *There exists a left ideal J in a C^* -algebra, and a surjective complete isometry $A \rightarrow J$ which is both a homomorphism (i.e. multiplicative), and a triple morphism.*

Proof. It is trivial that (ii) implies (i), and that (iii) implies (ii) (since we can choose a nonnegative r.c.a.i. in J). In fact it is not too hard to see that (iii) implies (ii) for *any* r.c.a.i. for A , but we will not need this.

Finally, given (i), we observe that by hypothesis, A is a triple envelope of A . Let $\mathcal{E}$ be the ‘left C^* -algebra’ of this triple system, and let $\langle \cdot | \cdot \rangle : A \times A \rightarrow \mathcal{E}$ be the associated sesquilinear inner product. By [9] Theorem 2.3, the map $\psi(x) = \lim_\alpha \langle x | e_\alpha \rangle$ is a well defined completely isometric triple morphism of A onto a left ideal J of $\mathcal{E}$, and $\psi = \psi \circ Id_A$ is also a homomorphism. $\square$

Remarks. (1). There are various rather trivial ‘TRO’ characterizations of left ideals in C^* -algebras. For example, one can characterize left ideals in C^* -algebras with right identities as the TRO’s Z which possess an element e of norm 1 such that $xe = x$ for all $x \in Z$. This is equivalent to $xe^* = x$ for all $x \in Z$. A similar condition with a net does the general case. Indeed a similar idea gives a rather trivial characterization of left ideals in C^* -algebras as the ‘abstract triple systems’ which are also operator algebras with a r.c.a.i. if one adds algebraic compatibility conditions such as (ii) above, between the operator algebra product and the triple product. Our result seems a little deeper than this. The only compatibility condition we seem to need between the operator algebra and the triple product is that the induced matrix norms are the same.

(2). A slight modification of this result also gives a characterization of C^* -algebras, by replacing ‘r.c.a.i.’ by ‘c.a.i.’. We are grateful to Bernie Russo for pointing out a recent paper [18] which gives such a characterization, but without needing the matrix norms.

(3) It would be interesting if, in the spirit of [25], one could give a purely linear characterization of left ideals in C^* -algebras. There is such a result in [8], but it makes reference to the containing C^* -algebra in the hypotheses.

Corollary 2.15 allows us to generalize the main result of [7] (see also [24]) to algebras with one-sided c.a.i.:

Theorem 2.19. *Let A be an operator algebra with r.c.a.i., which has a predual operator space. Then A has a right identity e of norm 1. Also A is a ‘dual operator algebra’, which means that the product on A is separately weak* continuous, and there exists a completely isometric homomorphism, which is also a homeomorphism with respect to the weak* topologies, of A onto a σ -weakly (i.e. weak*-) closed subalgebra B of some $B(H)$.*

Proof. The first assertion appears in [7] Theorem 2.5 (indeed for this part we only need a predual Banach space). From [7] Theorem 3.2, $\mathcal{M}_\ell(A)$ is a dual operator algebra. We saw in 2.13 and 2.15 that $\lambda : A \rightarrow \mathcal{M}_\ell(A)$ is a completely isometric homomorphism onto a left ideal

of $\mathcal{M}_\ell(A)$. Hence $\lambda(A) = \mathcal{M}_\ell(A)\lambda(e)$. Thus $\lambda(A)$ is a weak* closed subalgebra of $\mathcal{M}_\ell(A)$, and so $B = \lambda(A)$ is a dual operator algebra. If we take a bounded net $\lambda(a_i) \rightarrow \lambda(a)$ weak* in $\lambda(A)$, then by definition of the weak* topology on $\mathcal{M}_\ell(A)$ from [7] 3.2, $a_i e = a_i \rightarrow a e = a$ weak* in A . Thus λ^{-1} is weak* continuous, so that by Krein-Smulian (c.f. [7] Lemma 1.5) λ is weak* continuous. $\square$

The following ‘1-sided version’ of Sakai’s theorem may be known (certainly most of it is contained in a result of Zettl [33] (recently given a new proof in [15], which one may greatly shorten in the case where one has an operator space predual [7])).

Theorem 2.20. *Let J be a left ideal in a C^* -algebra, and suppose that J possesses a Banach space predual. Then $M(JJ^*)$ is a W^* -algebra containing J as a weak*-closed principal left ideal.*

Proof. By the aforementioned result of Zettl, $M(JJ^*)$ is a W^* -algebra and J is a dual operator space. By 2.19, J has a right identity e , so that $J = JJ^*e = M(JJ^*)e$. $\square$

Results such as 2.14 and 2.16 are useful for deducing results about general operator algebras with r.c.a.i., from results about left ideals in C^* -algebras. Here is a sample application of this ‘transference principle’ (other examples will be given later):

Corollary 2.21. *Let A be an operator algebra with a right contractive approximate identity, and also a right identity. Then A has a right identity of norm 1, which is the limit in norm of the r.c.a.i..*

Proof. If $\{e_\alpha\}$ is the r.c.a.i. for A , then $\{j(e_\alpha)\}$ is a r.c.a.i. for the left ideal envelope $\mathfrak{J}_e(A)$. Similarly $\mathfrak{J}_e(A)$ and A have a common right identity. Hence by 2.6, our r.c.a.i. converges in norm. $\square$

3. THE BANACH-STONE THEOREM

We prove several stages, or cases, of this theorem, which asserts essentially that linear surjective complete isometries between left ideals of C^* -algebras (resp. between operator algebras with r.c.a.i.), are characterized as a composition of a translation by a partial isometry u , and a surjective completely isometric homomorphism onto another right ideal (resp. operator algebra with r.c.a.i.) which is a translate of one of the original ideals (resp. algebras) by u^* . To see that the ‘translate by a partial isometry’ is not artificial, consider an infinite dimensional Hilbert space H and S the shift operator. Set $I = B(H)$ and $J = B(H)S$. These ideals are clearly linearly completely isometric, but there is no homomorphism of I onto J (since J has no 2-sided identity). This example shows that the following theorem (which comprises Case (1)) is best possible:

Theorem 3.1. *Let I and J be principal left ideals in C^* -algebras A and B , i.e. $I = Ae$ and $J = Bf$, say, for orthogonal projections e, f in I, J respectively. Suppose also that $\varphi : I \rightarrow J$ is a linear surjective complete isometry. Then there exists a partial isometry u in B with initial projection f , and a completely isometric surjective ideal homomorphism $\pi : I \rightarrow J_1$ such that $\varphi = \pi(\cdot)u$ and $\pi = \varphi(\cdot)u^*$. Here $J_1 = Bu^* = Ju^* = Buu^* \subset B$ is another left ideal of B with right identity uu^* of norm 1.*

Conversely, if J is a left ideal of a C^ -algebra B , and if u is a partial isometry in B with initial projection a right identity for J of norm 1, then $Ju^* = Bu^* = Buu^*$ is a left ideal J_1 of B with right identity uu^* of norm 1, and J_1 is linearly completely isometrically isomorphic to J via right multiplication by u^* . Hence the composition of right multiplication by u^* , with any completely isometric surjective homomorphism $I \rightarrow J_1$, is a linear completely isometric isomorphism $I \rightarrow J$.*

Proof. Recall from the introduction that a completely isometric surjection between TRO's is a triple morphism. Hence φ is a triple isomorphism. Therefore if $u = \varphi(e)$ then it is easy to check that $\pi(\cdot) = \varphi(\cdot)u^*$ is a homomorphism onto Ju^* . Similar considerations show that $p = uu^*$ is an idempotent, which is an orthogonal projection since it is selfadjoint. Thus u is a partial isometry. We claim that $u^*u = f$. To see this note that u^*u is an orthogonal projection, and that for any $\varphi(x) \in J$ we have $\varphi(x)u^*u = \varphi(xe) = \varphi(x)$, using the definition of a triple morphism. Thus $fu^*u = f$. On the other hand, $u^*uf = u^*u$ since $u \in Bf$. Hence $f = u^*u$. Also, $Ju^* = Bfu^* = Bu^*uu^* = Bu^*$. Defining J_1 to be this last space we see that it is clearly a left ideal of B , and J_1 contains uu^* , which is indeed a right identity of norm 1 for J_1 since u is a partial isometry. Thus $J_1 = Buu^*$ too.

Since $\pi(\cdot) = \varphi(\cdot)u^*$ we obtain $\pi(\cdot)u = \varphi(\cdot)u^*u = \varphi(\cdot)f = \varphi(\cdot)$. It follows from this too that π is a complete isometry, and therefore also a triple morphism. Thus π is a completely isometric ideal homomorphism.

Conversely, if J, B, u are as stated, then $J = Bu^*u$ so that $Ju^* = Bu^*$ which is also a right ideal of B . Clearly the last space equals Buu^* since $Buu^* \subset Bu^* = Bu^*uu^* \subset Buu^*$. The remainder of the converse direction is left to the reader. $\square$

Thus we see that central to the Banach-Stone theorem in our 1-sided case, is a certain Murray - von Neumann equivalence of projections.

Note in the 'unital case' of the above theorem, i.e. the case that $\varphi(e) = f$, then $u = u^* = f$, $J_1 = J$, and φ itself is a homomorphism. Conversely, if φ is a homomorphism, then necessarily $\varphi(e) = f$, as is easy to see from 2.7.

Having thoroughly analyzed the Banach-Stone theorem in Case (1), we now move to Case (2). Here we look at linear completely isometric isomorphisms $\varphi : A \rightarrow B$ between operator algebras with a right identity of norm 1. In the assertions in the first paragraph of the statement of the theorem, and in the proofs of these assertions, we regard A and B as having been identified with subalgebras of $\mathfrak{J}_e(A)$ and $\mathfrak{J}_e(B)$ respectively (see 2.14). Thus mention of the 'canonical Shilov embedding homomorphisms' j have been suppressed, and all products and adjoints in that paragraph are taken in the containing C^* -algebra $\mathcal{E}(B) = \mathfrak{J}_e(B)\mathfrak{J}_e(B)^*$.

Theorem 3.2. *(Banach-Stone for operator algebras with right identities.) Suppose that $\varphi : A \rightarrow B$ is a surjective linear completely isometric isomorphism between operator algebras with a right identity of norm 1. Then there exists a partial isometry $u \in \mathfrak{J}_e(B)$ (indeed in B) with initial projection the right identity of B , such that the subspace $B' = Bu^*$ of $\mathcal{E}(B)$ is a subalgebra (and consequently an operator algebra) with a right identity uu^* of norm 1; and there exists a completely isometric surjective homomorphism $\pi : A \rightarrow B'$, such that $\varphi = \pi(\cdot)u$ and $\pi = \varphi(\cdot)u^*$. Also, $u^*B \subset B$.*

Conversely, suppose we are given a partial isometry u on a Hilbert space H , such that u lies in a subalgebra $B \subset B(H)$, such that the initial projection of u is a right identity of B , and such that $u^*B \subset B$. Then $B' = Bu^*$ is an operator algebra with right identity uu^* of norm 1, and B' is linearly completely isometrically isomorphic to B via right multiplication by u . Thus the composition of right multiplication by u^* , with any completely isometric surjective homomorphism $A \rightarrow B'$, is a linear completely isometric isomorphism $A \rightarrow B$.

We remark that in the language of [9], u and u^* are in $LM(B)$.

Proof. Suppose that $\varphi : A \rightarrow B$ is a linear completely isometric isomorphism, and extend φ to a linear completely isometric isomorphism $\bar{\varphi} : \mathfrak{J}_e(A) \rightarrow \mathfrak{J}_e(B)$ (such extension exists by Hamana theory ([21] or [6] Appendix A)). By Theorem 2.14, $\mathfrak{J}_e(A)$ is a left ideal of the C^* -algebra $\mathcal{E}(A)$, and $\mathfrak{J}_e(A)$ has right identity e . Similar assertions hold for $\mathfrak{J}_e(B)$. Thus by the proof of 3.1, if $u = \varphi(e) = \bar{\varphi}(e)$ then u is a partial isometry in B , with $u^* \in B^* \subset \mathcal{T}(B)^* \subset \mathcal{E}(B)$, whose initial projection is f , and $\pi = \bar{\varphi}(\cdot)u^*$ is a completely isometric surjective homomorphism $\mathfrak{J}_e(A) \rightarrow \mathfrak{J}_e(B)u^*$. The restriction of π to A maps onto the subalgebra Bu^* of $\mathcal{E}(B)$. Since u is a partial isometry, uu^* is indeed a right identity of Bu^* . Finally, since $Bu^*Bu^* \subset Bu^*$, post multiplying by u gives $Bu^*B \subset B$, so that $u^*B = u^*uu^*B = fu^*B \subset Bu^*B \subset B$.

Conversely, given u as stated, then since $u^*B \subset B$ we have that Bu^* is a subalgebra of $\mathcal{E}(B)$ with right identity uu^* . The remainder of the converse direction is obvious. $\square$

One can prove further that $\mathfrak{J}_e(B') = \mathfrak{J}_e(B)u^*$, and that $\mathcal{E}(B') = \mathcal{E}(B)$, but we omit the details.

Corollary 3.3. *Suppose that $\varphi : A \rightarrow B$ is a surjective linear completely isometric isomorphism between operator algebras with right identities e and f of norm 1. Then φ is a homomorphism if and only if $\varphi(e) = f$.*

Proof. The one direction follows from 2.7, the other by noting that if we follow the proof of 3.2, then $\bar{\varphi}(e) = f$, so that $\bar{\varphi}$ is a homomorphism by the remarks after 3.1. $\square$

Corollary 3.4. *Suppose that A is an operator algebra with a right identity of norm 1, and suppose that A has another product $m : A \times A \rightarrow A$ with respect to which A is completely isometrically isomorphic to an operator algebra with a right identity of norm 1. Then there is a partial isometry $u \in \mathfrak{J}_e(A)$ (and indeed in A) such that $m(x, y) = xu^*y$ for all $x, y \in A$. Indeed u is the right identity for m , and u^*u is the right identity for the first product.*

This result also has corollaries of the type given in [3]; we leave the details to the reader.

We now turn to Case (3) of the Banach-Stone theorem:

Theorem 3.5. *(The Banach-Stone theorem for left ideals in C^* -algebras.) Consider a surjective linear complete isometry $\varphi : I \rightarrow J$ between arbitrary left ideals of C^* -algebras. Let $\mathcal{E} = JJ^*$, and let $\mathcal{M}$ be the von Neumann algebra $(JJ^*)^{**}$. Then there exists another left ideal J_1 of $\mathcal{E}$, with $J_1J_1^* = \mathcal{E}$, and a surjective completely isometric ideal homomorphism $\pi : I \rightarrow J_1$. Moreover there exists a partial isometry $u \in \mathcal{M}$ such that the initial projection of u is the right identity of J^{**} (indeed of $RM(J)$ - see §4), and such that $J_1 = Ju^*$, $J = J_1u$, and such that $\varphi = \pi(\cdot)u$, and $\pi = \varphi(\cdot)u^*$.*

There clearly exists a (tidier) converse to this theorem, as in Cases (1) and (2).

Proof. Consider the second dual $\varphi^{**} : I^{**} \rightarrow J^{**} \subset \mathcal{M}$, and now we are back in Case (1). For if I is a left ideal of a C^* -algebra A , then I^{**} is a left ideal of A^{**} , but now I^{**} has a right identity e of norm 1, namely e is a weak*-accumulation point of the r.c.a.i. of I (by [13] 28.7). Thus by Case (1) we have that $u = \varphi^{**}(e)$ is a partial isometry in $J^{**} \subset \mathcal{M}$, and the initial projection of u is the matching right identity of J^{**} . Moreover $\pi = \varphi^{**}(\cdot)u^*$ is a completely isometric homomorphism and so on. Restricting π to I gives a completely isometric homomorphism π' onto the subalgebra $J_0 = Ju^*$ of $\mathcal{M}$, and φ is the composition of π' with a right translation by u . Moreover, π' is easily seen to be a triple morphism:

$$\pi'(x)\pi'(y)^*\pi'(z) = \varphi^{**}(\hat{x})u^*u\varphi^{**}(\hat{y})^*\varphi^{**}(\hat{z})u = \varphi(x)\varphi(y)^*\varphi(z)u = \varphi(xy^*z)u = \pi'(xy^*z)$$

for $x, y, z \in I$. Thus π' is a completely isometric ideal homomorphism. Therefore by Lemma 2.10, π' is the restriction of a surjective 1-1 *-homomorphism $II^* \rightarrow J_0J_0^*$. Thus $J_0J_0^* = Ju^*uJ^* = \mathcal{E}$ contains J_0 as a left ideal; or to be more precise, $\hat{\mathcal{E}}$ contains J_0 . Thus we may regard π' as a completely isometric homomorphism $\pi : I \rightarrow J_1$ onto a right ideal J_1 of $\mathcal{E}$ (note $\hat{J}_1 = J_0$). The rest is clear. $\square$

Finally, Case (4) of the Banach-Stone theorem, i.e. the case of a surjective linear complete isometry between arbitrary operator algebras with r.c.a.i.. Again it is clear that by passing to the second dual and using Case (2) in the way we tackled Case (3) using Case (1), *or* using Case (3) in the way we tackled Case (2) using Case (1), will give a correct theorem resembling Theorems 3.1, 3.2, and 3.5. We leave the details to the reader. This should also be linked to the multiplier algebras in [9].

Corollary 3.6. *Let $\varphi : A \rightarrow B$ be a surjective linear complete isometry between left ideals of C^* -algebras, or between operator algebras with r.c.a.i.. Then φ is a homomorphism if and only if there exists a r.c.a.i. $\{e_\alpha\}$ for A such that $\{\varphi(e_\alpha)\}$ is a r.c.a.i. for B .*

Proof. If the latter condition holds then $\varphi^{**} : A^{**} \rightarrow B^{**}$ is a surjective linear complete isometry. Let E be a weak* limit point of $\{e_\alpha\}$ in A^{**} , and since φ^{**} is weak*-continuous, $\varphi^{**}(E)$ is a weak* limit point of $\{\varphi(e_\alpha)\}$. So we are in the situation of Corollary 3.3 by 2.8, so that φ^{**} and consequently φ is a homomorphism. The converse direction is easier. $\square$

4. $LM(A)$ FOR AN ALGEBRA WITH LEFT CONTRACTIVE APPROXIMATE IDENTITY

In this section we develop the ‘left multiplier operator algebra’ $LM(A)$ of an operator algebra with l.c.a.i. (note that of course $RM(A)$ for an operator algebra with r.c.a.i. will have the almost identical, ‘other-handed version’, of this theory). On the other hand the left multiplier operator algebra of an operator algebra with r.c.a.i. turns out to have a quite different theory, which is studied in the sequel [9], and which we will not mention again in the present paper.

Actually, it appears at the present time as if there may be two classes of ‘good candidates’ for $LM(A)$ if A has l.c.a.i.. We will only really investigate the first of these classes in this section, because in this case we can show that all the candidates in this class coincide, and we get a convincing theory paralleling the known theory in many ways. The one drawback

of this left multiplier algebra is that it does not contain the algebra itself in general; but this is no surprise or mental obstacle to anyone who has looked at the ‘multiplier’ or ‘centralizer’ theory of nonunital Banach algebras. The second class of ‘candidates’ consist of algebras which do contain A , but then of necessity one loses the important property that every nontrivial multiplier may be viewed as a nontrivial map $A \rightarrow A$, and this introduces some problems. Indeed this second class of ‘candidates’ seems to be a rather unruly zoo; although there seems to be a ‘best’ or ‘most canonical’ candidate in this class, which we call the ‘big left multiplier algebra’ $BLM(A)$. We will briefly discuss these matters further at the end of this section (see also Remark 3 after 4.2).

Thus we restrict our attention to the first class of candidates. Since this follows closely the essentially known theory for the case of a two-sided c.a.i., we will try to be brief. As a historical note, the latter theory was begun in [31], and also independently developed around that time from a different angle, mostly by Paulsen [28], as a tool for our project on Morita equivalence [10], although this latter material was not circulated or published. Some other facts about this case of an operator algebra with a two-sided c.a.i. have arisen in several of the authors papers over the years, for example [4, 6]. However since these facts were not particularly profound, followed mostly from the ideas of [31] and [28], and did not play a crucial role, we did not give a complete development in these places, essentially leaving the needed details to the reader. The majority of these facts are contained in the following result, and its proof. In view of this result, if A is an operator algebra with l.c.a.i. then we will write $LM(A)$ for any of the completely isometrically isomorphic algebras of the theorem, and $\mu_A : A \rightarrow LM(A)$ for the canonical map (see also Remark 2 below). To understand this result better it is good to have a simple example in mind, such as $A = R_n$ (the subalgebra of M_n supported on the first row).

Theorem 4.1. *Let A be an operator algebra with l.c.a.i.. Then the following operator algebras are all completely isometrically isomorphic:*

- (1) $\{x \in A^{**} : x\hat{A} \subset \hat{A}\} / \ker q$ where q is the canonical homomorphism into $CB(A)$,
- (2) $\mathcal{M}_\ell(A)$,
- (3) $CB_A(A)$ (right module maps).

Moreover $CB_A(A) = B_A(A)$ isometrically. If A has a two-sided c.a.i., then $\ker q = (0)$ in (1), and also the above algebras also are completely isometrically isomorphic to:

- (4) the ‘maximum essential left multiplier extension of A ’.

If A satisfies condition $(\mathcal{L})$ of 2.2 (for example if A has a left identity of norm 1, or a two-sided c.a.i., or if A is a right ideal of a C^* -algebra), then the algebras above are completely isometrically isomorphic to

- (5) $\{T \in B(H) : T\pi(A) \subset \pi(A)\}$, for any completely $\mathcal{L}$ -isometric nondegenerate representation of A (see definition after 2.12),
- (6) $LM(B)$ where $B = \mathcal{L}(A)$ (see 2.2),
- (7) $\{x \in B^{**} : x\hat{A} \subset \hat{A}\} \subset A^{**}$, where $B = \mathcal{L}(A)$.

To avoid distracting from other issues here, we will not explain the term ‘essential left multiplier extension’ used in (4) until we reach it in the proof below.

Proof. We first observe that for *any* operator algebra A there are natural completely contractive homomorphisms

$$\{x \in A^{**} : x\hat{A} \subset \hat{A}\} \rightarrow \mathcal{M}_\ell(A) \rightarrow CB(A).$$

Let us write σ for the first homomorphism, and θ for the second. From the ‘left handed variant’ of 2.15, the image of θ lies in $CB_A(A)$. The fact that $CB_A(A) = B_A(A)$ isometrically follows more or less immediately from the relation $T(a) = \lim_\alpha T(e_\alpha)a$. Next note that given $S \in CB_A(A)$, then one may let F be a weak* accumulation point of $S(e_\alpha)$ in A^{**} , for the l.c.a.i. $\{e_\alpha\}$ for A . Clearly $\|F\| \leq \|S\|$. For $a \in A$, we have

$$S(a) = \lim_\alpha S(e_\alpha a) = \lim_\alpha S(e_\alpha)a = Fa.$$

Hence $q(F) = S$ where $q = \theta \circ \sigma$. Thus q is a quotient map, and similarly it is a complete quotient map. Thus σ is also a complete quotient map, and $\ker \sigma = \ker q$ since θ is 1-1. This proves the completely isometric isomorphism between (1) and (3), and also between (1) and (2). Thus $\mathcal{M}_\ell(A) \cong CB_A(A)$ completely isometrically, which also shows that $CB_A(A)$ is a unital operator algebra (or this fact may be proved directly).

Now suppose that A has property $(\mathcal{L})$, and set $B = \mathcal{L}(A)$ as in 2.2. Then $B^{**} \subset A^{**}$. Examining the proof of (1) = (3) above, we see easily that the terms $S(e_\alpha)$ actually lie in B . Hence the F there lies in $\{x \in B^{**} : x\hat{A} \subset \hat{A}\}$. Thus the map q mentioned above, restricted to the last set, is a complete quotient map too. Therefore it is a complete isometry if we can show that it is 1-1. To see this suppose that F is in the set in (7) and $q(F) = 0$. Then $F\hat{e}_\alpha = 0$. This implies that $F = 0$, using the fact that a weak* limit point of the $\hat{e}_\alpha$ is a 2-sided identity for B^{**} , and the fact that the multiplication in a dual operator algebra is separately weak* continuous. Thus we have that (3) = (7) completely isometrically. Note too that if A is an operator algebra with 2-sided c.a.i. then this shows that $\ker q = (0)$ in (1). Note that if F is in the set in (7), then $FB \subset B$ quite clearly. Conversely if $FB \subset B$ then for $a \in A$ we have $Fa = \lim Fe_\alpha a \in A$ since $Fe_\alpha a \in Ba \subset A$. This shows that (6) = (7).

Returning to (4), we need a definition of an ‘essential left multiplier extension’ of an operator algebra A with c.a.i.. For the purposes of this theorem we will define this to be a pair (B, π) consisting of an operator algebra B with identity of norm 1, together with a completely isometric homomorphism $\pi : A \rightarrow B$, such that $\pi(A)$ is a left ideal of B , and such that the natural map $B \rightarrow CB(A)$ is completely isometric. There is a natural ordering, and corresponding notion of equivalence, of ‘left multiplier extensions’ of A , which we will not bother to spell out. However it is clear that since the just mentioned ‘natural map’ maps into $CB_A(A)$, the algebras in (1)-(3) are the biggest essential left multiplier extensions.

Finally to prove that (5) = (6), we may w.l.o.g., by the definition after 2.12 and the last assertion of 2.12, assume that $B = A$ is an operator algebra with 2-sided c.a.i., and that $\pi : A \rightarrow B(H)$ is a nondegenerate completely isometric homomorphism. This case is well known; briefly, one way to see it is as follows. Noting that the algebra in (5) is then an essential left multiplier extension of A , we see that the algebra in (5) is completely isometrically contained as a subalgebra of $CB_A(A)$. Conversely, if $R \in B_A(A)$, we obtain a related map $T \in B(H)$ which may be defined by $T\pi(a)\zeta = \pi(Ta)\zeta$, for $a \in A, \zeta \in H$. Another way to see this quickly is using the well known fact that in this case, $H \cong A \hat{\otimes}_A H$.

We omit the simple details, which as we said at the beginning of this section, are essentially well known to experts. $\square$

Corollary 4.2. *Let A be an operator algebra with left identity e of norm 1. Then $LM(A) = Ae$, which is a unital subalgebra of A . It is also a unital subalgebra of $\mathcal{E}(A)$, and $\mathcal{E}(A)$ is a unital C^* -algebra.*

Proof. In this case $\mathcal{L}(A) = Ae$ which is a unital algebra. Thus the first result here follows from (6) of the previous theorem. We saw in 2.14 that $J = \mathfrak{J}_e(A)$ is a right ideal of a C^* -algebra, and J has a left identity e . Thus $\mathcal{E}(A) = JJ^*$ has e as a 2-sided identity. Finally $Ae \subset JJ^* = \mathcal{E}(A)$. $\square$

Remark 1. There is a condition similar to (4) in 4.1 which is equivalent to (1)-(3) in general. However it was sufficiently more complicated to offset the benefits of mentioning it in more detail.

Remark 2. It is very important that any kind of multiplier algebra D of an algebra A should not merely be regarded as an algebra, but rather as a *pair* (D, μ) consisting also of a homomorphism $\mu : A \rightarrow D$. Sometimes we write μ_A to indicate the dependence on A . Saying that two such algebras (D, μ) and (D', μ') are the same as multiplier algebras must entail a completely isometric surjective homomorphism $\theta : D \rightarrow D'$ such that $\theta \circ \mu = \mu'$. In this case we say that D and D' are A -isomorphic. Thus in each of the seven equivalent formulations of the previous theorem, we need to have in mind also what the map μ is in each case. In (1) it is the map $a \mapsto \hat{a} + \ker q$; in (2) and (3) it is essentially the regular representation λ ; in (5) it is π ; in (6) it is the natural left representation of A on its left ideal $\mathcal{L}(A)$; and in (7) it is $a \mapsto \hat{a}E$, where E is as in the remark before 2.12. All these maps are completely contractive homomorphisms. One then needs to check that these seven algebras are all A -isomorphic. We leave these assertions to the reader who wishes to be more careful.

Remark 3. Suppose that A is an operator algebra with l.c.a.i., and that $\pi : A \rightarrow B(H)$ is a completely isometric representation. Define $LM(\pi) = \{T \in B(H) : T\pi(A) \subset \pi(A)\}$, the left idealizer of $\pi(A)$ in $B(H)$. Then it is straightforward to exhibit a completely contractive homomorphism $\sigma : LM(\pi) \rightarrow LM(A) = CB_A(A)$. Conversely, given $T \in CB_A(A)$, taking a weak operator limit point S of $\pi(T(e_\alpha))$ gives $S \in LM(\pi)$. This is really saying that $LM(A) \cong LM(\pi)/\ker \sigma$ completely isometrically isomorphically. One may view this observation as an attempt to remove the use of property $(\mathcal{L})$ in (5). Perhaps an investigation of this quotient might be tied to Sarason's semi-invariant subspace technique (see [1] for example). It is interesting to note that if π is the usual representation of R_2 , then $LM(\pi)$ is a 3-dimensional operator algebra (this was pointed out to me by M. Kaneda). Note that $LM(\pi)$ is highly dependent on π , to see this consider R_2 again; the natural representation π has $LM(\pi)$ 3-dimensional. However, if $\sigma = \pi \oplus \epsilon$, where ϵ is the projection onto the 1-1 coordinate, then $LM(\sigma)$ is strictly larger. It would be interesting to see if there is a nonrestrictive condition under which one obtains 'independence from the particular π used'.

We now turn to the notion which in the C^* -algebra literature is referred to as 'essential homomorphisms' or sometimes 'nondegenerate homomorphisms'. For our purpose we shall

use the name ‘ A -nondegenerate morphism’. For us this shall mean a completely contractive homomorphism $\pi : A \rightarrow LM(B)$ satisfying the following equivalent conditions:

Theorem 4.3. *Let A and B be two operator algebras with l.c.a.i.’s, and let $\pi : A \rightarrow LM(B)$ be a completely contractive homomorphism. The following are equivalent:*

- (i) *There exists a l.c.a.i. $\{e_\alpha\}$ for A such that $\pi(e_\alpha)b \rightarrow b$ for all $b \in B$,*
- (ii) *For every l.c.a.i. $\{e_\alpha\}$ for A , we have $\pi(e_\alpha)b \rightarrow b$,*
- (iii) *B is a nondegenerate left A -module via π ,*
- (iv) *Any $b \in B$ may be written $b = \pi(a)b'$ for some $a \in A, b' \in B$.*

If these conditions hold, there exists a completely contractive unital homomorphism $\hat{\pi} : LM(A) \rightarrow LM(B)$ such that $\hat{\pi} \circ \mu_A = \mu_B$. Moreover $\hat{\pi}$ has the property that $\hat{\pi}(x)(\pi(a)b) = \pi(xa)b$ for $x \in LM(A), a \in A, b \in B$; and it is the unique such homomorphism with this property. Finally $\hat{\pi}$ is completely isometric if π is completely isometric.

Proof. Clearly (i) implies that the span of terms $\pi(a)b$ is dense in B , which is what we mean by nondegenerate. So (i) implies (iii). Clearly (iii) implies (ii), and (ii) implies (i), and (iv) implies (iii). That (iii) implies (iv) follows from [26] §5.2.

If these conditions hold, view $LM(A)$ and $LM(B)$ as in (3) of 4.1, and note the formula for μ_A in the previous remark. We may follow the proof of Theorem 6.2 in [4] (which the author proved inspired by an argument of van Daele mentioned there). The main difference is that we ignore the element e mentioned there, which we can get away with by taking d there to be the l.c.a.i. from A . One also needs to use [26] 5.2.2, and the matricial version of it, in order to show that $\hat{\pi} : LM(A) \rightarrow CB_B(B)$, and that it is a complete contraction. The remaining problem is the last assertion. Supposing that π is completely isometric, we have

$$\|\hat{\pi}(T)\|_{cb} \geq \|[\hat{\pi}(T)(\pi(a_{ij})b_{kl})]\| = \|[\pi(Ta_{ij})b_{kl}]\|$$

providing that $\|[a_{ij}]\|, \|[b_{kl}]\| \leq 1$. Taking the supremum over all such $[b_{kl}] \in M_m(B)$, gives that

$$\|\hat{\pi}(T)\|_{cb} \geq \|[\pi(Ta_{ij})]\| = \|[Ta_{ij}]\|.$$

Taking the supremum over all such $[a_{ij}] \in M_n(A)$ gives that $\|\hat{\pi}(T)\|_{cb} \geq \|T\|_{cb}$. So $\hat{\pi}$ is isometric, and similarly it is completely isometric. $\square$

We note that the canonical map $\mu_A : A \rightarrow LM(A)$ is an A -nondegenerate morphism. Note too that if $\pi : A \rightarrow B$ is a completely contractive homomorphism between two operator algebras with l.c.a.i.’s, then we can say that π is A -nondegenerate, if $\mu_B \circ \pi$ is A -nondegenerate. This happens if and only if π takes a l.c.a.i. for A to a l.c.a.i. for B , or equivalently that B is a nondegenerate A -module via π . Thus the inclusion map from a closed subalgebra A of an operator algebra B is A -nondegenerate if A contains a l.c.a.i. for B . In this case we have

Corollary 4.4. *Let A be a closed subalgebra of an operator algebra B , and suppose that A contains a l.c.a.i. for B . Then $LM(A) \hookrightarrow LM(B)$ completely isometrically as a subalgebra.*

We end with the promised remarks about a second class of candidates for the ‘left multiplier operator algebra’ of an operator algebra A with l.c.a.i.. For a candidate (B, ν) in this class one would like to have the property that $\nu : A \rightarrow B$ is a completely isometric homomorphism,

such that $B\nu(A) \subset \nu(A)$. Unfortunately then one must lose the useful ‘essential’ condition (i.e. that $x\nu(A) = 0$ implies $x = 0$). Thus for many A (such as our standard example R_n) a subset of B plays no role in any action on A , and this fact does not seem to bode well for ‘uniqueness’ properties of such multiplier algebras. Indeed this takes us out of the classical ‘multiplier’/‘centralizer’ framework from Banach algebra theory ([26] §1.2 for example), where a multiplier which annihilates A must be the zero multiplier. Also one cannot hope for conditions like (1)-(3) of 4.1, it appears.

With these cautions it seems nonetheless that there is a ‘most interesting candidate’ in this class. Consider the injective C^* -algebra B and homomorphism j of (the other-handed version of) Theorem 2.16, and define $BLM(A) = \{x \in B : xj(A) \subset j(A)\}$. That is, $BLM(A)$ is the left idealizer of $j(A)$ in B . For example, $BLM(R_2)$ is the upper triangular 2×2 matrix algebra. Since $I_{11} = eBe \subset B = I_{22}$ in the language of (the other-handed version of) 2.16, it may be proved by the multiplication theorem in [11] and 1.3 there, that $BLM(A)$ is the ‘biggest’ unital operator algebra containing A completely isometrically as a subalgebra, in the sense that it contains a completely contractive image of every other such algebra C . It may not contain C itself though (consider $A = R_2$, and see Remark 3 above).

Other candidate definitions for a left multiplier algebra containing A might look like the $LM(\pi)$ algebras in Remark 3 after 4.2. One might hope that there is a suitable and not too strong condition on π there so that these algebras are independent of the particular π , but this seems unlikely at present. It seems as if one could present large numbers of further such candidates, which at this point seem unrelated in any way.

REFERENCES

- [1] W. B. Arveson, *Subalgebras of C^* -algebras*, Acta Math. **123** (1969), 141-224.
- [2] W. B. Arveson, *Subalgebras of C^* -algebras II*, Acta Math. **128** (1972), 271-308.
- [3] D. P. Blecher, *Commutativity in operator algebras*, Proc. Amer. Math. Soc. **109** (1990), 709-715.
- [4] D. P. Blecher, *A generalization of Hilbert modules*, J. Funct. An. **136** (1996), 365-421.
- [5] D. P. Blecher, *Some general theory of operator algebras and their modules*, in *Operator algebras and applications*, A. Katavolos (editor), NATO ASIC, Vol. 495, Kluwer, Dordrecht, 1997.
- [6] D. P. Blecher, *The Shilov boundary of an operator space, and the characterization theorems*, J. Funct. An. **182** (2001), 280-343.
- [7] D. P. Blecher, *Multipliers and dual operator algebras*, J. Funct. An. **183** (2001), 498-525.
- [8] D. P. Blecher, E. G. Effros and V. Zarikian, *One-sided M -ideals and multipliers in operator spaces I*, Preprint (2000). To appear *Pacific J. Math.*
- [9] D. P. Blecher and M. Kaneda, *The ideal envelope of an operator algebra*, Preprint.
- [10] D. P. Blecher, P. S. Muhly and V. I. Paulsen, *Categories of operator modules - Morita equivalence and projective modules*, *Memoirs of the American Mathematical Society* Vol. 143, number 681 (Jan, 2000).
- [11] D. P. Blecher and V. I. Paulsen, *Multipliers of operator spaces, and the injective envelope*, Pacific J. Math. **200** (2001), 1-17.
- [12] D. P. Blecher, Z. J. Ruan, and A. M. Sinclair, *A characterization of operator algebras*, J. Funct. An. **89** (1990), 188-201.
- [13] F. F. Bonsall and J. Duncan, *Complete normed algebras*, Springer-Verlag, New York-Heidelberg (1973).
- [14] J. B. Conway, *A course in functional analysis*, Graduate Texts in Mathematics, Springer-Verlag, 1985.
- [15] E. G. Effros, N. Ozawa and Z. J. Ruan, *On injectivity and nuclearity for operator spaces*, Duke Math. J. To appear.
- [16] E. G. Effros and Z. J. Ruan, *On non-self-adjoint operator algebras*, Proc. Amer. Math. Soc. **110** (1990), 915-922.

- [17] E. G. Effros and Z. J. Ruan, *Operator Spaces*, Oxford University Press (2000).
- [18] K. El Amin, A. M. Campoy, and A. Rodriguez-Palacios, *A holomorphic characterization of C^* - and JB^* -algebras*, Manuscripta Math. **104** (2001), 467-478.
- [19] P. A. Fillmore, *A users guide to operator algebras*, Canadian Math. Soc. series, Wiley-Interscience (1996).
- [20] M. Hamana, *Injective envelopes of operator systems*, Publ. R.I.M.S. Kyoto Univ. **15** (1979), 773-785.
- [21] M. Hamana, *Triple envelopes and Silov boundaries of operator spaces*, Math. J. Toyama University **22** (1999), 77-93.
- [22] R. V. Kadison, *Isometries of operator algebras*, Ann. of Math. **54** (1951), 325-338.
- [23] M. Kaneda and V. I. Paulsen, *Characterizations of essential ideals as operator modules over C^* -algebras*, Preprint (2000).
- [24] C. Le Merdy, *An operator space characterization of dual operator algebras*, Amer. J. Math., **121** (1999), 55-63.
- [25] M. Neal and B. Russo, *A holomorphic characterization of ternary rings of operators*, Preprint, 2001.
- [26] T. W. Palmer, *Banach Algebras and the general theory of $*$ -algebras Vol. I*, Encyclopedia of mathematics and its applications Vol. 49, Cambridge University Press (1994).
- [27] V. I. Paulsen, *Completely bounded maps and dilations*, Pitman Research Notes in Math., Longman, London, 1986.
- [28] V. I. Paulsen, *Unpublished notes on multiplier algebras*, (circa 1992).
- [29] G. Pederson, *C^* -algebras and their automorphism groups*, Academic Press (1979).
- [30] G. Pisier, *Introduction to the theory of operator spaces*, to appear.
- [31] Y-t. Poon and Z. J. Ruan, *Operator algebras with contractive approximate identities*, Canadian J. Math. **46**(1994), 397-414.
- [32] N. E. Wegge-Olsen, *K -theory and C^* -algebras*, Oxford Univ. Press (1993).
- [33] H. Zettl, *A characterization of ternary rings of operators*, Adv. Math. **48** (1983), 117-143.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HOUSTON, 4800 CALHOUN, HOUSTON, TX 77204-3008, U.S.A.

E-mail address: dblecher@math.uh.edu